



# **MARKSCHEME**

**May 1999**

**MATHEMATICS**

**Higher Level**

**Paper 1**

1. By the remainder theorem,


$$f(-1) = 6 - 11 - 22 - a + 6 \quad (M1)$$

$$= -20 \quad (M1)$$

$$\Leftrightarrow a = -1 \quad (A2)$$

$$\text{Answer: } a = -1 \quad (C4)$$

2. Using a tree diagram,



$$p(\text{BG or GB}) = \left(\frac{3}{9} \times \frac{4}{8}\right) + \left(\frac{4}{9} \times \frac{3}{8}\right) \quad (M1)(M1)$$

$$= \frac{1}{6} + \frac{1}{6} \quad (A1)$$

$$= \frac{1}{3} \quad (A1)$$

$$\text{OR } p(\text{BG or GB}) = 2 \times \frac{4}{9} \times \frac{3}{8} \quad (M1)(M1)$$

$$= \frac{1}{3} \quad (A2)$$

$$\text{Answer: } \frac{1}{3} \quad (C4)$$

3. Let  $a$  be the first term and  $d$  be the common difference, then

$$a + d = 7 \text{ and } S_4 = \frac{4}{2}(2a + 3d) = 12 \quad (M1)$$

$$\Rightarrow \begin{cases} a + d = 7 \\ 4a + 6d = 12 \end{cases} \quad (M1)$$

$$\Rightarrow a = 15, d = -8 \quad (A2)$$

Answer:  $a = 15$  (C2)

$d = -8$  (C2)

4. Substituting gives,

$$2(2\lambda + 4) + 3(-\lambda - 2) - (3\lambda + 2) = 2 \quad (M1)$$

$$\Leftrightarrow 4\lambda + 8 - 3\lambda - 6 - 3\lambda - 2 = 2 \quad (M1)$$

$$\Leftrightarrow \begin{aligned} -2\lambda &= 2 \\ \lambda &= -1 \end{aligned} \quad (A1)$$

Intersection is  $(2, -1, -1)$  (A1)

Answer: Intersection is  $(2, -1, -1)$  (C4)

5.  $(1-i)z = 1-3i$

$$\Leftrightarrow z = \frac{1-3i}{1-i} \quad (M1)$$

$$\Leftrightarrow z = \frac{1-3i}{1-i} \times \frac{1+i}{1+i} \quad (M1)$$

$$\Leftrightarrow z = 2-i \quad (A2)$$

OR Let  $z = x + iy$

$$(1-i)(x+iy) = 1-3i \quad (M1)$$

$$x + y - i(x - y) = 1 - 3i$$

$$\begin{cases} x + y = 1 \\ x - y = 3 \end{cases} \quad (M1)$$

$$\Rightarrow x = 2, y = -1 \quad (A2)$$

Answer:  $x = 2$  (C2)

$y = -1$  (C2)

Note: Award (C4) for  $z = 2 - i$

6. The system of equations will not have a unique solution if the determinant of the matrix representing the equations is equal to zero.

Therefore,  $\begin{vmatrix} 4 & -1 & 2 \\ 2 & 3 & 0 \\ 1 & -2 & a \end{vmatrix} = 0$  (M1)


$\Leftrightarrow 4 \times 3a + 2a + 2 \times (-4 - 3) = 0$  (M1)

$\Leftrightarrow 14a = 14$  (M1)

$a = 1$  (A1)

Answer:  $a = 1$  (C4)

7.



(A2)(A2)

Notes: The graph of  $y_2$  is  $y_1$  shifted  $k$  units to the right.

Award (A2) for the correct graph.

Award (A1) for indicating each point of intersection with the  $x$ -axis i.e.  $(m+k, 0)$  and  $(n+k, 0)$

Answer: See graph (C4)

Note: Award (C4) if the graph of  $y_2$  is drawn correctly and correctly labelled with  $m+k$  and  $n+k$ .

8. Using a tree diagram,



(M2)

Let  $p(S)$  be the probability that the pupil speaks Spanish.

Let  $p(A)$  be the probability that the pupil is Argentine.

Then, from diagram,

$$p(S|A) = \frac{12}{15} \quad (A1)$$


$$= \frac{4}{5} \quad (A1)$$

OR  $p(S|A) = \frac{p(S \cap A)}{p(A)} \quad (M1)$

$$= \frac{12}{21} / \frac{15}{21} \quad (M1)(A1)$$

$$= \frac{4}{5} \quad (A1)$$

OR



(M2)

$$p(S|A) = \frac{12}{15} \quad (A1)$$

$$= \frac{4}{5} \quad (A1)$$

Answer:  $p(S|A) = \frac{4}{5} \quad (C4)$

9. By implicit differentiation,

$$\frac{d}{dx}(2x^2 - 3y^2 = 2) \Rightarrow 4x - 6y \frac{dy}{dx} = 0 \quad (M1)$$

$$\Leftrightarrow \frac{dy}{dx} = \frac{4x}{6y} \quad (A1)$$

$$\text{When } x = 5, 50 - 3y^2 = 2$$

$$\Leftrightarrow y^2 = 16 \quad (M1)$$

$$\Leftrightarrow y = \pm 4$$

$$\text{Therefore } \frac{dy}{dx} = \pm \frac{5}{6} \quad (A1)$$

Note: This can be done explicitly

$$\text{Answers: } \frac{dy}{dx} = \pm \frac{5}{6} \quad (C2)(C2)$$

10. (a) A perpendicular vector can be found from the vector product

$$\vec{OP} \times \vec{OQ} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -3 & 2 \\ -2 & 1 & -1 \end{vmatrix} = \vec{i} - 3\vec{j} - 5\vec{k} \quad (M1)(A1)$$

$$(b) \text{ Area } \triangle OPQ = \frac{1}{2} \left| \vec{OP} \right| \left| \vec{OQ} \right| \sin \theta, \text{ where } \theta \text{ is the angle between } \vec{OP} \text{ and } \vec{OQ} \quad (M1)$$

$$\begin{aligned} &= \frac{1}{2} \left| \vec{OP} \times \vec{OQ} \right| \\ &= \frac{\sqrt{35}}{2} \quad (A1) \end{aligned}$$

$$\text{Answers: (a) } \vec{i} - 3\vec{j} - 5\vec{k} \text{ (or any multiple)} \quad (C2)$$

$$(b) \frac{\sqrt{35}}{2} \quad (C2)$$

11. Given  $y = \arccos(1 - 2x^2)$

then  $\frac{dy}{dx} = \frac{-1}{(1 - (1 - 2x^2)^2)^{1/2}} \times -4x$  (M1)

$$\frac{dy}{dx} = \frac{4x}{(1 - (1 - 4x^2 + 4x^4))^{1/2}} \quad (M1)$$

$$\frac{dy}{dx} = \frac{4x}{(4x^2 - 4x^4)^{1/2}} \quad (A2)$$

OR

$$\cos y = 1 - 2x^2 \quad (M1)$$

$$-\sin y \frac{dy}{dx} = -4x$$

$$\frac{dy}{dx} = \frac{-4x}{-\sin y} = \frac{4x}{\sqrt{1 - (1 - 2x^2)^2}} \quad (M1)$$

$$\frac{dy}{dx} = \frac{4x}{\sqrt{4x^2 - 4x^4}} \quad (A2)$$

Answer:  $\frac{dy}{dx} = \frac{4x}{(4x^2 - 4x^4)^{1/2}}$  or  $\frac{dy}{dx} = \frac{4x}{\sqrt{4x^2 - 4x^4}}$  (C4)

12. Let  $f(x) = ax^2 + bx + c$  where  $a = 1$ ,  $b = (2 - k)$  and  $c = k^2$ . Then for  $a > 0$ ,  $f(x) > 0$  for all real values of  $x$  if and only if

$$b^2 - 4ac < 0 \quad (M1)$$


$$\Leftrightarrow (2 - k)^2 - 4k^2 < 0 \quad (A1)$$

$$\Leftrightarrow 4 - 4k + k^2 - 4k^2 < 0$$

$$\Leftrightarrow 3k^2 + 4k - 4 > 0$$

$$\Leftrightarrow (3k - 2)(k + 2) > 0 \quad (M1)$$

$$\Leftrightarrow k > \frac{2}{3}, k < -2 \quad (A1)$$



**Answer:**  $k < -2$ , or  $k > \frac{2}{3}$  (C2)(C2)

13. Let the volume of the solid of revolution be  $V$ .

$$V = \pi \int_0^a ((ax + 2)^2 - (x^2 + 2)^2) dx \quad (M1)$$

$$= \pi \int_0^a (a^2 x^2 + 4ax + 4 - x^4 - 4x^3 - 4) dx \quad (M1)$$

$$= \pi \left[ \frac{1}{3} a^2 x^3 + 2ax^2 - \frac{1}{5} x^5 - \frac{4}{3} x^3 \right]_0^a \quad (M1)$$

$$= \pi \left( \frac{2}{15} a^5 + \frac{2}{3} a^3 \right) \text{ units}^3 \quad (A1)$$

$$= \frac{2a^3\pi}{15} (a^2 + 5)$$

**Note:** The last line is not required

**Answer:**  $V = \frac{2a^3\pi}{15} (a^2 + 5)$  or equivalent (C4)



14. Let  $u = \frac{1}{2}x + 1 \Leftrightarrow x = 2(u-1) \Rightarrow \frac{dx}{du} = 2$

Then  $\int x \left( \frac{1}{2}x + 1 \right)^{1/2} dx = \int 2(u-1) \times u^{1/2} \times 2 du$  (M1)

$$= 4 \int (u^{3/2} - u^{1/2}) du$$
 (A1)

$$= 4 \left[ \frac{2}{5} u^{5/2} - \frac{2}{3} u^{3/2} \right] + C$$
 (M1)

$$= 4 \left[ \frac{2}{5} \left( \frac{1}{2}x + 1 \right)^{5/2} - \frac{2}{3} \left( \frac{1}{2}x + 1 \right)^{3/2} \right] + C$$
 (A1)

$$= \frac{8}{15} \left( \frac{1}{2}x + 1 \right)^{3/2} \left( \frac{3}{2}x - 2 \right) + C$$

Note: The last line is not required

Answer:  $4 \left[ \frac{2}{5} \left( \frac{1}{2}x + 1 \right)^{5/2} - \frac{2}{3} \left( \frac{1}{2}x + 1 \right)^{3/2} \right] + C$  or  $\frac{8}{15} \left( \frac{1}{2}x + 1 \right)^{3/2} \left( \frac{3}{2}x - 2 \right) + C$  (C4)

15. The locus defined by  $|z - 4 - 3i| = |z - 2 + i|$  is the perpendicular bisector of the line joining the points  $4 + 3i$  and  $2 - i$  in the complex plane.

Correct diagram (see below).

(M2)

(A2)

OR

Let  $z = x + yi$  then,  $|z - 4 - 3i| = |z - 2 + i|$

$$\Leftrightarrow (x-4)^2 + (y-3)^2 = (x-2)^2 + (y+1)^2$$
 (M1)


$$\Leftrightarrow -8x - 6y + 25 = -4x + 2y + 5$$

$$\Leftrightarrow x + 2y = 5$$
 (M1)

Therefore the equation of the locus is  $x + 2y = 5$ .

Correct diagram (see below).

(A2)



Note: Award (A1) each for any two of the following (maximum of [2 marks]):  
Perpendicular line, midpoint (3, 1), y-intercept, x-intercept, gradient.

Answer: Correct locus (see diagram)

(C4)

16. Given  $(1+x)^5(1+ax)^6 = 1+bx+10x^2+\dots+a^6x^{11}$   
 $\Leftrightarrow (1+5x+10x^2+\dots)(1+6ax+15a^2x^2+\dots) = 1+bx+10x^2+\dots+a^6x^{11}$  (M1)  
 $\Leftrightarrow 1+(6a+5)x+(15a^2+30a+10)x^2+\dots = 1+bx+10x^2+\dots+a^6x^{11}$  (M1)

$$\Leftrightarrow \begin{aligned} 6a+5 &= b & \textcircled{1} \\ 15a^2+30a+10 &= 10 & \textcircled{2} \end{aligned}$$

$$\begin{aligned} \textcircled{2} \quad \Rightarrow \quad 15a^2+30a &= 0 & (M1) \\ \Leftrightarrow 15a(a+2) &= 0 \\ \Rightarrow a &= -2 \end{aligned}$$

$$\text{Substitute into } \textcircled{1} \quad b = -7 \quad (A1)$$

**Note:**  $a \neq 0$  since  $a \in \mathbb{Z}^*$

$$\text{Answers: } \begin{aligned} a &= -2 & (C2) \\ b &= -7 & (C2) \end{aligned}$$

17. Let  $X$  be the number of counters the player receives in return.

$$E(X) = \sum p(x) \times x = 9 \quad (M1)$$

$$\Leftrightarrow \left(\frac{1}{2} \times 4\right) + \left(\frac{1}{5} \times 5\right) + \left(\frac{1}{5} \times 15\right) + \left(\frac{1}{10} \times n\right) = 9 \quad (M1)(A1)$$

$$\Leftrightarrow \frac{1}{10}n = 3$$

$$\Leftrightarrow n = 30 \quad (A1)$$

$$\text{Answer: } n = 30 \quad (C4)$$

18. Let  $\Phi(z) = 0.017$   
 then  $\Phi(-z) = 1 - 0.017 = 0.983$  (M1)  
 $z = -2.12$  (A1)

$$\text{But } z = \frac{x - \mu}{\sigma} = \frac{1 - 1.02}{\sigma} \text{ where } x = 1 \text{ kg}$$

$$\text{Therefore } \frac{1 - 1.02}{\sigma} = -2.12 \quad (M1)$$

$$\Leftrightarrow \sigma = 0.00943 \text{ kg} = 9.4 \text{ g (to the nearest } 0.1 \text{ g)} \quad (A1)$$

$$\text{Answer: } \sigma = 9.4 \text{ g (or equivalent)} \quad (C4)$$

19. (a) Given  $\frac{dv}{dt} = -kv$

$$\Leftrightarrow \int \frac{dv}{v} = -k \int dt$$

$$\Leftrightarrow \ln v = -kt + C \quad (M1)$$

$$\Leftrightarrow v = Ae^{-kt} \quad (A = e^C)$$

At  $t = 0$ ,  $v = v_0 \Rightarrow A = v_0$

$$\Leftrightarrow v = v_0 e^{-kt} \quad (A1)$$

(b) Put  $v = \frac{v_0}{2}$

then  $\frac{v_0}{2} = v_0 e^{-kt}$  (M1)

$$\Leftrightarrow \frac{1}{2} = e^{-kt}$$

$$\Leftrightarrow \ln \frac{1}{2} = -kt$$

$$\Leftrightarrow t = \frac{\ln 2}{k} \quad (A1)$$

Note: Accept equivalent forms, e.g.  $t = \frac{\ln \frac{1}{2}}{-k}$

Answers: (a)  $v = v_0 e^{-kt}$  (C2)

(b)  $t = \frac{\ln 2}{k}$  (C2)

20. Given  $v = \frac{(3s+2)}{(2s-1)}$

then acceleration  $a = \frac{dv}{dt} = \frac{dv}{ds} \times \frac{ds}{dt} = \frac{dv}{ds} \times v$  (M1)

therefore  $a = \frac{3(2s-1) - 2(3s+2)}{(2s-1)^2} \times \frac{(3s+2)}{(2s-1)}$  (M1)

$$\Leftrightarrow a = \frac{-7(3s+2)}{(2s-1)^3} \quad (M1)$$

therefore when  $s = 2$ ,  $a = \frac{-56}{27}$  (A1)

Answer: acceleration =  $-\frac{56}{27}$  (C4)